

COMPOSITION :

Each 1 gm contains :
Amoxicillin (as amoxicillin trihydrate).....800 mg



AMOXY 800 - 1KG

ANTIBIOTIC SOLUBLE POWDER FOR ORAL ADMINISTRATION

INDICATIONS:

It is freely water - soluble, suitable for all types of potable water.

Amoxicillin is a hemi-synthetic and a broad-spectrum penicillin, acting against bacteria's GRAM-positive and GRAM-negative. Levels of increasing resistance have been observed concerning E.coli, Salmonella typhimurium and Pasteurella haemolytica.

Calves: Respiratory infections caused by Pasteurella spp. and Salmonella dublin.

Gastrointestinal infections caused by E.coli.

Infections caused by Salmonella dublin.

Poultry: Respiratory infections caused by E.coli, Haemophilus spp, Pasteurella spp and Salmonella spp. gastrointestinal infections caused by Salmonella spp.

CONTRAINDICATIONS:

Do not administer to animals with hypersensitivity to penicillins and cephalosporins, to animals with serious renal disorders and other animal species not listed at the indications.

INTERACTION WITH OTHER MEDICINAL PRODUCT:

Do not administer simultaneously with effective bacteriostatic antibiotics such as tetracyclines, etc

WITHDRAWAL PERIOD:

Calves: Meat: 2 days, Milk: 2 days.

Poultry: Meat: 1 day, Egg: 0 day.

DOSAGE AND ADMINISTRATION:

Calves: 50 g per 100 liters of milk substitute or potable water twice a day, for 5 consecutive days (10 mg amoxicillin per kg of body weight /12 hours)

Poultry: 15 mg amoxicillin trihydrate(equivalent to 13.1 mg amoxicillin) per kg body weight per day (corresponding to 19 mg of AMOXY product/ kg b.w/day).

The total period of treatment should be for 3 days or in severe cases for 5 days.

The medicated water must be replaced every 12 hours. The medicated milk replacer must be administered within the next 6 hours.

It is recommended to use gloves and mask during the administration of the product, especially in case of known hypersensitivity to penicillin.

Keep it In original pack, Store in a dry, dark place below 25°C.

USED FOR

Calves & Poultry

