



## WITHDRAWAL PERIODS:

Cattle: Meat - 10 days • Milk - 35 days  
Sheep and Goats: Meat - 10 days • Milk - 35 days  
Do not use in animals producing milk for human consumption during or within the withdrawal period.

# WORMVET PLUS

**DESCRIPTION:** A broad-spectrum oral anthelmintic combining Albendazole and Levamisole HCl, supported with key trace minerals (Cu, Zn, Se, Co) to aid immune and metabolic recovery after parasitic stress.

This dual-action formula effectively eliminates mature and immature gastrointestinal and lungworms, as well as tapeworms and adult liver flukes in cattle and sheep. Opti-Gent Spray helps control and reduce local bacterial infections sensitive to gentamicin, including: Gram-positive species (*Staphylococcus aureus*, *Streptococcus* spp., *Corynebacterium* spp.) and Gram-negative species (*E. coli*, *Proteus*, *Salmonella*, *Pseudomonas*, *Moraxella*, etc.). It is particularly useful for bovine keratoconjunctivitis (pink-eye) caused by *Moraxella bovis*, as well as superficial wounds, cuts, abrasions, and umbilical infections.

**INDICATIONS:** Indicated for the treatment and prevention of susceptible gastrointestinal and lungworm infestations, and for the removal of adult liver flukes and tapeworms in cattle and sheep. Cattle: Effective against *Ostertagia* (*Teladorsagia*), *Haemonchus*, *Trichostrongylus*, *Bunostomum*, *Nematodirus*, *Oesophagostomum*, *Cooperia*, and *Strongyloides* spp., including inhibited larvae. Also active against lungworms (*Dictyocaulus viviparus*), tapeworms (*Moniezia* spp.), and adult liver flukes (*Fasciola hepatica*). Sheep: Active against *Teladorsagia* (*Ostertagia*), *Haemonchus*, *Trichostrongylus*, *Nematodirus* (including *N. battus*), *Chabertia*, and *Oesophagostomum* spp., including inhibited larvae, as well as lungworms (*Dictyocaulus filaria*), tapeworms (*Moniezia* spp.), and adult liver fluke (*Fasciola hepatica*).

Albendazole also exhibits ovicidal properties, destroying the eggs of flukes and nematodes to reduce reinfection and pasture contamination. The trace mineral blend aids post-treatment recovery, supporting enzyme systems, red-blood-cell production, and fertility improvement after deworming.

## COMPOSITION:

EACH 1 ML CONTAIN:	
Albendazole	23.8mg
Levamisole HCl	37.5mg
Selenium	0.5mg
Cobalt	0.3mg
Copper	2.1mg
Zinc	0.6mg



## DOSAGE AND ADMINISTRATION:

**For oral administration only.** Do not mix with other veterinary products or feed additives in the same dosing equipment.

Each mL of WORMVET PLUS Suspension delivers approximately 4.75 mg/kg Albendazole and 7.5 mg/kg Levamisole HCl when administered at the standard rate.

**CATTLE:** Worm dose: For the control of gastrointestinal and pulmonary nematodes, tapeworms, and to reduce fluke and roundworm egg output.

Dosage: 1 mL per 5 kg body weight.

Fluke and worm dose: For additional treatment against adult liver fluke (chronic fascioliasis) in cattle.

Dosage: 1 mL per 5 kg body weight.

**SHEEP:** Worm dose: For the control of roundworms, lungworms, tapeworms, and to reduce fluke and worm egg shedding.

Dosage: 1 mL per 5 kg body weight.

Fluke and worm dose: For additional treatment of adult liver fluke (chronic fascioliasis) in sheep.

Dosage: 1 mL per 5 kg body weight.

**STORE BELOW 30°C in the original container**

## USED FOR

Cattle & Sheep